

Methodological concerns in evaluating water exchange combined with distal attachments for adenoma detection

Selim Coşkun^{a,b}, Shota Hoshika^{b,c},
Masahiro Banno^{b,d}

Koseiren Murakami General Hospital, Niigata; Scientific Research Works Peer Support Group, Osaka; Funabashi Orthopaedic Sports Medicine & Joint Centre, Chiba; Seichiryō Hospital, Nagoya, Japan

We read with interest the study by Leung *et al* regarding water exchange with distal attachments [1]. While the authors conclude that this combination optimizes the right-sided adenoma detection rate (ADR), several methodological issues suggest that these findings should be interpreted with caution.

First, the primary endpoint (overall ADR) did not reach statistical significance ($P=0.05$). The authors' suggestion that the non-significant result for Endocuff Vision ($P=0.057$) was merely a Type II error is speculative. As emphasized by Amrhein *et al* [2], in rigorous clinical trials, results exceeding the $P<0.05$ threshold should not be framed as positive trends to justify clinical efficacy.

Second, the reported benefit in the right colon was a non-prespecified secondary outcome. According to the trial registration (NCT03566615), right-sided ADR was not a predefined endpoint. Highlighting this specific finding after a negative primary result raises concerns about selective reporting and the inflation of Type I error through multiple testing.

Furthermore, the 13-min mean withdrawal time significantly exceeds the 6-8 min guideline recommendation and the 9-min clinical average [3]. This prolonged inspection is likely to create a "ceiling effect" [4], where meticulous observation outweighs the incremental benefit of distal attachments, limiting the results' generalizability to routine practice.

In summary, given the *post-hoc* nature of the significant findings, the borderline statistical results, and the non-standard withdrawal times, the conclusion that these modalities should be routinely used to optimize right-sided ADR is premature. A more conservative, hypothesis-generating tone would be more appropriate.

References

1. Leung F, Cadoni S, Chen L, et al. A prospective international multisite randomized controlled trial of water exchange with and without distal cap(s) in adenoma detection. *Ann Gastroenterol* 2026;**39**:104-113.
2. Amrhein V, Greenland S, McShane B. Scientists rise up against statistical significance. *Nature* 2019;**567**:305-307.
3. Garborg KK. How many minutes should we take for withdrawal - guideline recommendations and upcoming evidence. *Best Pract Res Clin Gastroenterol* 2026;**80**:102019.
4. Shaukat A, Rector TS, Church TR, et al. Longer withdrawal time is associated with a reduced incidence of interval cancer after screening colonoscopy. *Gastroenterology* 2015;**149**:952-957.

^aDivision of Gastroenterology, Koseiren Murakami General Hospital, Niigata, Japan (Selim Coşkun); ^bScientific Research Works Peer Support Group (SRWS-PSG), Osaka, Japan (Selim Coşkun, Shota Hoshika, Masahiro Banno); ^cShoulder & Elbow Service, Funabashi Orthopaedic Sports Medicine & Joint Center, Chiba, Japan (Shota Hoshika); ^dDepartment of Psychiatry, Seichiryō Hospital, Nagoya, Japan (Masahiro Banno)

Conflict of Interest: None, Funding: None

Correspondence to: Selim Coşkun MD, Division of Gastroenterology, Koseiren Murakami General Hospital, 5-8-1 Midoricho, Murakami-shi, Niigata, 958-8533, Japan, e-mail: 18selimcos@gmail.com

Received 7 February 2026; accepted 23 February 2026; published online 23 April 2026

DOI: <https://doi.org/10.20524/aog.2026.1063>

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms